

FERTILIZATION PROGRAM: WATERMELON











VIGORTEM S

4-5 kg/ha



Fertilizer with non-mycorrhizal microorganisms

TRICOBEST 1 kg/ha · 3 applications

Root system biostimulant

First irrigation: 1-2 kg/ha · 10 days after the first application: 2-3 kg/ha

VIGORTEM S

Metabolic biostimulant, improves

flowering and fruit setting

Flowering biostimulant

Amino acid biostimulant, enhances fruit enlargement, color and sheen

LIDEX PREMIUM

 $2.5 \text{ kg/ha} \cdot 3 \text{ applic.}$: with the beehive's introduction, 10 days before and 7 days after

MOLIBOR EXTRA 150-200 g/100 L

> **MAXITEM** 5-7 L/ha

Note: The above fertilization program it's a general recommendation based on the experience and studies of Quimicas Meristem. It should serve as a guide, having to be adapted to each individual case depending on both local conditions and crop characteristics.



Product certified for eco-friendly agriculture A Foliar application



Root application

FERTILIZATION PROGRAM: WATERMELON PRODUCT USE RECOMMENDATIONS



TRICOBEST

TRICOBEST is a high quality and high soluble concentrate of humic acids, enriched with *Trichoderma harzanium* (strain IABTH01). It has a notable biostimulant effect, enhancing root development and crop vigor. The strain helps solubilize phosphorus and chelate other nutrients, increasing plan availability. Humic acids improve physical and chemical soil properties. 2-3 applications with an interval of 7-10 days.



VIGORTEM S

VIGORTEM S combines a high phosphorus content, humic and fulvic acids, seaweed extract and aminoacids. VIGORTEM S promotes root growth and vigor ensuring crop establishment, and due to humic and fulvic acids, improves physical and chemical soil properties.



LIDEX PREMIUM

LIDEX PREMIUM is a metabolic processes biostimulant that promote crop development and improves flowering and fruit setting, optimizing crop productivity.





MOLIBOR EXTRA

MOLIBOR EXTRA stimulates the early onset of pollen and improves its quality, resulting more viable flowers. MOLIBOR EXTRA reduces flower abscission, enhancing the setting process. Apply 7 days before and 2 days after beehives introduction.



MAXITEM

MAXITEM combines fulvic, that improves soil structure acids with amino acids that enhance nutrient absorption and movement through the plant. MAXITEM enhances crop vigor and promotes fruit size, color, and sheen.

